(12)

EUROPEAN PATENT APPLICATION published in accordance with Art. 158(3) EPC

- (43) Date of publication: 26.09.2001 Bulletin 2001/39
- (21) Application number: 99974093.9
- (22) Date of filing: 30.09.1999

- (51) Int Cl.7: C12N 15/82, A01H 5/00
- (86) International application number: PCT/JP99/05386
- (87) International publication number: WO 01/25459 (12.04.2001 Gazette 2001/15)
- (84) Designated Contracting States:

 AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU

 MC NL PT SE
- (71) Applicant: Japan Tobacco Inc. Tokyo 105-8422 (JP)
- (72) Inventors:
 - KURAYA, Yoshiki, Japan Tobacco Inc. lwata-gun, Sizuoka 438-0802 (JP)

- KOMARI, Toshihiko, Japan Tobacco Inc. Iwata-gun, Sizuoka 438-0802 (JP)
- HIEI, Yukoh, Japan Tobacco Inc. Iwata-gun, Sizuoka 438-0802 (JP)
- (74) Representative:
 Reinhard Skuhra Weise & Partner
 Postfach 44 01 51
 80750 München (DE)

(54) VECTORS FOR TRANSFORMING PLANTS

(57) Vectors for transforming plants with the use of agrobacteria which have been modified so as to elevate the possibility of the recognition of the border sequences of the vectors by vir proteins of the agrobacteria, thereby lowering the possibility of the transfer of DNAs other than T-DNA into plant chromosomes. More particularly, the above-vectors are those to be used in transforming

plants which have right and left border sequences which can be recognized by the vir proteins of the agrobacteria, a T-DNA sequence which is located between these border sequences and into which a gene to be transferred into plants can be inserted, and a replication origin enabling the replication of the vectors in bacteria, characterized by having a plural number of left border sequences.

Description

TECHNICAL FIELD

[0001] The present invention relates to plant transformation vectors, more particularly, vectors useful in *Agrobacte-rium*-mediated plant transformation. The invention further relates to a method of transforming plants using the vectors. The invention is particularly useful for generating transgenic plants that may be taken as food.

BACKGROUND ART

10

[0002] It has long been known that *Agrobacterium* (*Agrobacterium tumefaciens*), is a soil bacterium which causes Crown gall disease in many dicotyledonous plants. In the ninety-seventies, it was found that the Ti plasmid of *Agrobacterium* is involved in pathogenicity and that T-DNA which is part of the Ti plasmid is integrated into the plant genome. It was later revealed that the T-DNA contained the hormone synthesis genes (cytokinins and auxins) necessary for crown gall tumorigenesis and that those genes, although derived from bacteria, are expressed in plants. A group of genes that are located in the virulence region (*Vir* region) of the Ti plasmid are necessary to the excision of T-DNA and its transfer to plants, and furthermore, the border sequences that are located on opposite ends of T-DNA are required for the excision, which are called the right border sequence and the left border sequence. *Agrobacterium rhizogenes*, another *Agrobacterium* species has a similar system involving the Ri plasmid.

[0003] Stated more specifically, the proteins produced on the basis of the genes located in the *vir* region (*vir* proteins) recognize the right and left border sequences to integrate the T-DNA located between the border sequences into plant genome. This function provided the basis for the transformation of plants with a foreign gene pre-inserted into T-DNA, thereby giving rise to the development of *Agrobacterium*-mediated plant transformation technology.

[0004] Most recently, however, several reports have appeared describing that, in certain kinds of plants, it is sometimes observed that T-DNA is not excised at the border sequences, and hence, T-DNA can be transferred into the plant chromosome together with a region adjacent to T-DNA (Ramanathan et al., Plant Molecular Biology 28, 1149-1154 (1995), and Kononov et al., Plant Journal 11, 945-957 (1997)). If a DNA element other than T-DNA is co-transferred, the resulting transgenic plants will be suspected of having unexpected characteristics, which could have a negative impact on public acceptance of food products made of transgenic plants. It is therefore desired to develop a method whereby it can be ensured that unnecessary non-T-DNA sequences of *Agrobacterium* will not transfer to plant chromosomes.

[0005] The inventors supposed that the *vir* proteins of *Agrobacterium* sometimes fail to recognize the border sequences and this may explain the reason why non-T-DNA is transferred into plant chromosome together with T-DNA. No vectors have yet been developed that can suppress or reduce the transfer of non-T-DNA segment with a view to solving said problem.

[0006] Based on the above supposition, the inventors have conducted intensive studies for creating vectors for use in *Agrobacterium*-mediated transformation. In order to reduce the probability that non-T-DNA element is transferred to plant chromosome the inventors modified the vector with a view to increasing the efficiency of the *vir* proteins of *Agrobacterium* to recognize the border sequence/s. As a result, it has been found that while two border sequences exist in the transformation vector, the probability of the integration of non-T-DNAs can be reduced by providing a plurality of left border sequences. The present invention has been accomplished on the basis of this finding.

SUMMARY OF THE INVENTION

[0007] The present invention provides a plant transformation vector based on the function of *Agrobacterium*, wherein the left border sequence has been modified such as to reduce the possibility of the integration of any non-T-DNA segment into plant chromosomes. More particularly, the invention provides a plant transformation vector comprising a right border sequence and a left border sequence that can be recognized by the *vir* proteins of *Agrobacterium*, a T-DNA region located between these border sequences and into which a gene to be introduced into the plant can be inserted, and a replication origin (*ori*) that enables replication of said vector in bacteria (e.g. *Agrobacterium* and bacteria for vector amplification), wherein said left border sequence has been modified such as to reduce the possibility of integration of any non-T-DNA sequence into plant chromosomes.

BRIEF DESCRIPTION OF THE DRAWINGS

[8000]

55

Fig. 1 shows vector pSLB0 used in the Example; this vector was prepared from pSB11 by modifying it to have

within th T-DNA region a cassette, for expr ssing the hygromycin resistanc gen under control of a ubiquitin promoter and a ubiquitin intron, then inserting a cassett for expressing the GUS gen containing a catalase intron under control of a ubiquitin promoter, at the site recognizable by restriction in nzyme Stul; and

Fig. 2 shows maps of the areas of vectors pSLB0, pSLB2 and pSLB3 in the new lighterhood of the left border sequence (which is hereunder sometimes referred to as LB). To prepare pSLB2 and pSLB3, a synthetic DNA fragment having two or three left border sequences was inserted into pSLB0 between LB and the GUS expressing cassette at the restriction site of Pvull. In pSLB2, two left border sequences were inserted by the synthetic DNA fragment (the inserted left border sequence/s is hereunder referred to as sLB) to give a total of three LBs, and in pSLB3, three sLBs were inserted by the synthetic DNA fragment to give a total of four LBs.

DETAILED DESCRIPTION OF THE INVENTION

(1) Vector preparation

5

10

35

40

45

50

[0009] The vector of the invention has the left border sequence modified to operate such as to reduce the possibility of the integration of an unnecessary DNA sequence/s, i.e. a non-T-DNA sequence/s into plant chromosomes. In a preferred embodiment, the invention provides the vectors having the left border sequence modified such as to comprise the placement of more than one DNA sequence that can be recognized by *vir* proteins (e.g. a known left border sequence).

[0010] DNA fragments comprising more than one left border sequence can be prepared by various known methods on the basis of known border sequences. For example, one can synthesize a single-stranded DNA molecule having the same sequence as a left border sequence which is contained in an available Ti plasmid, prepare the double-stranded DNA molecule from the single strand, and if necessary, link two or more of such double-stranded DNAs together. The obtained DNA fragment can be then inserted into a plant transformation vector at a suitable restriction site nearby up/downstream of the existing left border sequence located downstream of the T-DNA region. In this way, the vector of the invention can be easily constructed.

[0011] The plant transformation vector from which the vector of the invention can be prepared by modifying its left border sequence should at least have a right and a left border sequences that can be recognized by *vir* proteins, a T-DNA sequence located between the right and left border sequences and into which can be inserted a gene to be introduced into the plant, and a replication origin that can operate in bacteria for replication of the vector (e.g. *Escherichia coli*). The preferred plant transformation vector has a replication origin that can operate in *Agrobacterium*.

[0012] As long as these requirements are satisfied, various vectors can be modified on the left border sequence. For example, the various vectors used in the following plant transformation methods based on *Agrobacterium* can be modified:

(i) a small intermediate cloning vector having right and left border sequences and which has a foreign gene inserted into T-DNA, and an acceptor Ti plasmid having the *vir* region are subjected to homologous recombination to prepare a hybrid Ti plasmid vector, and the plant is infected with *Agrobacterium* containing the hybrid Ti plasmid vector; (ii) a foreign gene is inserted into the T-DNA region of a small Ti plasmid having no *vir* region (the plasmid is commonly called a mini plasmid or a micro Ti plasmid and is capable of replication in many bacteria) and the plasmid is introduced into *Agrobacterium* which harbors a plasmid having the *vir* region but no T-DNA, and the

(iii) a small intermediate cloning vector having a right and a left border sequences and which has an foreign gene inserted into T-DNA, and an acceptor Ti plasmid having a portion of the *vir* region (i.e., vir gene lacking a portion of the full length of the *vir* region) are subjected to homologous recombination to prepare a hybrid Ti plasmid vector, the hybrid Ti plasmid vector is introduced into *Agrobacterium* which has the *vir* region (full length) harboring but T-DNA deficient plasmid introduced in it, and the plant is infected with the *Agrobacterium* containing the two plasmids. The various vectors used in these methods can be modified on the left border sequence. Small Ti plasmids with a modification on the left border sequence are easy to handle in operations such as for modifying the foreign gene in T-DNA and hence are a preferred embodiment of the vector of the invention. Examples of such small Ti plasmids include pBI101 and pBI121 (both being available from CLONTECH), as well as pSB11 which was used in the Example to be described later.

[0013] The concept of the invention is applicable not only to the Ti plasmid but also to the Ri plasmid.

plant is infected with the Agrobacterium containing the two plasmids;

[0014] The vector of the invention may contain a marker gene in the T-DNA sequence that permits selection of the transformant, such as an antibiotic resistance gene or a luminescence gene. To be specific, commonly used marker genes may be employed in the usual manner and they include antibiotic resistance genes such as those conferring resistance to tetracycline, ampicillin, kanamycin, neomycin, hygromycin and spectinomycin, and luminescence genes

such as the luciferas gene, β -galactosidase, gre in fluorescence protein (GFP), β -lactamase and chloramphenicol acetyl transferase (CAT) genes. Besides thes genes, the vector may contain another marker gene outside of the T-DNA sequence, preferably downstream of the left border sequences. The marker gene placed in that position is useful for evaluating the effectiveness of the modified border sequences.

[0015] The term "replication origin" as used in this specification means a specific DNA region in which the replication reaction is initiated, commonly called *Ori*.

(2) Transformation

50

[0016] To use the vector of the invention, a foreign gene for the intended transformation is inserted into the T-DNA region. The foreign gene to be inserted usually contains a promoter that can operate in the host plant and the structural gene encoding the characteristic to be conferred to the plant linked downstream of the promoter. If necessary, more than one gene may be linked together and, in addition or alternatively, a sequence for enhancing the efficiency of expression may be interposed between the promoter and the downstream structural gene before insertion into the T-DNA region.

[0017] Before being introduced into a target plant, the vector of the invention which harbors the foreign gene is introduced into a bacterial of *Agrobacterium* species capable of infecting the plant (e.g. *Agrobacterium tumefaciens*). To this end, various methods well known to the skilled artisan can be employed. For example, the vector may be transferred into the *Agrobacterium* by conjugation; if possible, the *Agrobacterium* may be directly transformed with the vector of the invention containing the foreign gene.

[0018] Conventional techniques may be employed to infect the plant with the *Agrobacterium* containing the vector of the invention and they include, for example, wounding part of the plant body and infecting it with the bacterium, infecting the callus with the bacterium, co-cultivating the protoplast and the bacterium, and co-cultivating slices of the leaf tissue together with the bacterium. The transformed cells obtained by these methods can be selected by using the suitable selection marker/s or assaying if they express the intended characteristic. The transformed cells may further be differentiated by the prior art technology to yield a recombinant plant body.

[0019] In order to integrate that T-DNA containing the foreign gene into the chromosomal DNA in the plant, the *vir* region is necessary. The *vir* region may be supplied from the vector having the foreign gene or from a different vector.
[0020] The plant cells transformed with the vector of the invention may be differentiated by the prior art technology to yield a recombinant plant body. The transformed plant may be selected by using a suitable selection marker or assaying if it expresses the intended characteristic.

[0021] Whether a DNA sequence/s unnecessary for the intended transformation has been integrated into plant chromosomes or not can be determined by various methods well known to the skilled artisan. For example, oligonucleotide primers are synthesized on the basis of the vector DNA sequence/s outside the borders and with these primers, PCR is performed to analyze the chromosomal DNA sequences in the transformed plant. In the case of transformation with a vector that contains a marker gene outside the T-DNA sequence, analysis can be done by assaying if the marker gene has been expressed.

[0022] The vector of the invention is characterized by its function to reduce the possibility of the integration of a DNA sequence/s unnecessary for the intended transformation. The term "to reduce the possibility of the integration" means that, compared to the use of a vector which is not modified on the left border sequence, the frequency of the integration of the unnecessary DNA sequence/s into host chromosomes is low, or the length of the integrated unnecessary sequence is short, or there is no such integration and; in addition or alternatively, compared to the use of a vector which is not modified on the left border sequence, the frequency of unintended transformation is low, or unintended transformation is slight, or there is no occurrence of such transformation. The term "DNA sequence unnecessary for the intended transformation" means a portion or fragment of the DNA sequence located outside the T-DNA sequence in the vector (namely, non-T-DNA). It does not matter whether it is functional by itself or encodes a polypeptide or protein.

[0023] Various plants can be transformed by the transforming method of the invention and they include monocotyledonous plants such as maize, sorghum, triticale, barley, oats, rye, wheat, onion and rice, and dicotyledonous plants such as soybean, alfalfa, tobacco, rape, sunflower, potato, pepper and tomato. The method of the invention can reduce the possibility of the integration of a DNA sequence/s unnecessary for the intended transformation into plant chromosomes and the transgenic plants obtained by using the transformation method are less likely to have unexpected characteristics. Therefore, the method of the invention is suitable for transforming plants that can be taken as food by other organisms and regarding which there is particular concern about the possibility that the non-T-DNA sequence/s will transfer into plant chromosomes by *Agrobacterium*-mediated transformation. The method is most suitable for transforming monocotyledonous plants, in particular, rice.

[0024] Unless otherwise noted, the term "plant or plants" as used in the specification covers not only a plant body (individual) but also its s ed (germinated or immature), part (leaf, root, stem, flower, stamen, pistil or slices of these), culture c il, callus and protoplast.

Example

Example 1

5 (1) Preparing vectors

[0025] Plasmid vector pSB11 (Genbank Accession No. AB027256, Komari et al., Plant Journal 10, 165-174 (1966)) was modified to have within the T-DNA region a cassette for expressing the hygromycin resistance gene (HPT) by means of a ubiquitin promoter and a ubiquitin intron (Christensen et al., Plant Molecular Biology 18, 675-689 (1992) and into the plasmid, a cassette for expressing a catalase intron containing GUS gene (Ohta et al., Plant Cell Physiology 31, 805-813 (1990)) by means of a ubiquitin promoter was inserted at the site recognizable by restriction enzyme Stul. The thus prepared plasmid was designated pSLB0 (see Fig. 1 and SEQ ID:NO. 1). The nucleotide sequence of pSLB0 is shown as SEQ ID:NO. 1.

[0026] Then, based on the nucleotide sequence of Ti plasmid pTiAch5 (Genbank Accession No. K00548), a synthetic DNA containing a left border sequence (hereunder abbrevlated as LB) and the complementary synthetic DNA were prepared, they were annealed and processed to form blunt ends, and then used to prepare DNA fragments respectively having two and three LB sequences. The nucleotide sequences of the two synthetic DNAs are shown as SEQ ID:NO. 2 and SEQ ID:NO. 3. Each of the DNA fragments was inserted into pSLB0 between LB and the GUS expressing cassette at the site recognized by restriction enzyme Pvull; in this way, vectors were prepared that had more than one LB attached. The vector having two of the LB introduced by the synthetic DNA (which is hereunder sometimes referred to as sLB) to give a total of three LBs was designated pSLB2, and the vector having three sLBs introduced to give a total of four LBs was designated pSLB3 (for the maps of areas of pSLB0, pSLB2 and pSLB3 in the neighborhood of the synthetic LBs, see Fig. 2). Each of these three plasmids was introduced into Agrobacterium tumefaciens LBA4404 which already had plasmid vector pSB1 introduced (Genbank Accession No. AB027255, Komari et al., Plant Journal 10, 165-174 (1996)). They were subjected to the following tests.

(2) Transformation

[0027] Calli derived from the immature embryo of rice variety "Asanohikari" were transformed with LBA4404(pSLB0), LBA4404(pSLB2) and LBA4404(pSLB3) in accordance with the method of Hiel et al. (Hiel et al., Plant Journal 6, 271-282 (1994)).

- (3) Analysis of the expression of GUS gene in transformants
- [0028] Some leaves of the hygromycin-resistant plants obtained in Example 2 were stained with X-Gluc to check for the expression of the GUS gene. Seventeen out of the 340 plant individuals transformed with LBA4404(pSLB0) expressed the GUS gene, indicating that Agrobacterium derived DNA outside the border sequences was introduced into 5% of the plant individuals transformed with the conventional vector having only one LB. On the other hand, the number of plant individuals transformed with LBA4404(pSLB2) and LBA4404(pSLB3) and which expressed the GUS gene decreased with the increasing number of synthetic LBs (Table 1). This indicates that the integration of synthetic LBs into the vector decreased the likelihood for DNA beyond the left border sequence to be transferred to the plant.
 - (4) Analysis of genomic DNA in the individuals not expressing the GUS gene
- [0029] In some of the individuals that did not express the GUS gene in (3), DNA beyond the left border sequence may have been introduced into plant chromosomes but not far enough to the ubiquitin promoter for triggering the expression of the GUS gene. To verify this possibility in each group of plants that did not express the GUS gene, about 60 independent transformants were randomly chosen and genomic DNA was extracted and subjected to PCR analysis. The primers used in PCR analysis were so prepared as to permit amplification of the region extending from a location between the inherent LB and the synthetic LB to a location in the GUS gene. The sequences of the primers are shown as SEQ ID:NO. 4 and SEQ ID:NO. 5.

[0030] As a result of the PCR analysis, seven out of the 67 plant individuals (10.4%) transformed with LBA4404 (pSLB0) showed DNA amplification, revealing that when the conventional vector having only one LB was used, Agrobacterium derived DNA other than the desired T-DNA was integrated into chromosomes in the created transformants with a frequency of 10.4% of higher. In contrast, DNA amplification was found to take place in none of the plant individuals transformed with LBA4404(pSLB2) and LBA4404(pSLB3) that had synthetic LBs.

[0031] The results are shown in Tables 1 and 2.

Tabl 1.

	Analysis for the Expression of GUS Gen						
Vector	Number of transformants	Percentage of GUS expressing plants					
pSLB0	340	5.0					
pSLB2	327	1.2					
pSLB3	370	0.8					

Table 2.

Analysis of Genomic DNA							
No. of plants from which DNA was extracted	No. of plants in which DNA was amplified by PCR						
67	7						
5 5	0						
58	0						
	No. of plants from which DNA was extracted 67 55						

[0032] As can be seen from the above, the present invention decreased the integration of a DNA sequence/s outside the border sequences in plant chromosomes and made it possible to increase the efficiency of introducing only the intended T-DNA.

[0033] The foregoing description of the invention concerns primarily the use of two or more left border sequences, which may be derived from the same or different species of *Agrobacterium*. It should, however, be stressed that the fundamental concept of the invention lies in modifying the left border sequence in plant transformation vectors such that it can be recognized by *vir* proteins more efficiently to reduce the integration of any unnecessary non-T-DNA sequence to plant chromosomes. Therefore, the present invention embraces all vectors that have the modified left border sequence/s capable of achieving the same result. In addition to the examples described above, the modified left border sequences include the following: (1) those sequences which are derived from the sequence already existing in relevant plasmids by deletion, substitution or addition of one or more nucleotides in the existing left border sequence to be recognized by *vir* proteins more efficiently; (2) those sequences which are derived from the sequence already existing in the plasmid by deletion, substitution or addition of one or more bases in any sequence near the existing left border sequence to be recognized by *vir* proteins more efficiently; (3) those sequences which contain a plurality of any sequences that can be recognized by *vir* proteins; and (4) any combinations of (1) - (3).

SEQUENCE LISTING

<110> Japan	Tabacco	Inc
-------------	---------	-----

<120> Plant Transformation Vector

<130> YCT-441

<160> 5

5

10

15

30

35

40

45

50

55

20 <210> 1 <211> 12982

<212> DNA

25 (400) 1

.60 aagotigoat gootgoagig cagogigaco oggiogigoo cotototaga galaatgago attgcatgtc taagttataa aaaattacca catatttttt tigtcacact tgitigaagt 120 180 gcagtitale taictitata catalatita aactitacie taegaataat ataatetata gtactacaat aatatcagtg tiilagagaa tcatataaat gaacagttag acatggicta 240 aaggacaatt gagtattiig acaacaggac tctacagtii tatciittta gigigcaigt 300 360 gticicciti tittitigcaa atagciicac ciatataata ciicatccat titattagta caiccaitta gggittaggg ttaatggitt ttatagacta attititlag tacaictait 420 itaticiati itagecicia aattaagaaa actaaaacte tattitagii tittiatita 480 540 ataatttaga talaaaatag aalaaaataa agigactaaa aattaaacaa atacccilta 600 agaaattaaa aaaactaagg aaacattiit ciigiticga gtagataatg ccagccigit 660 aaacgccgic gacgagicia acggacacca accagcgaac cagcagcgic gcgicgggcc 720 aagcgaagca gacggcacgg catcicigic gcigcctcig gaccccicic gagagticcg ciccaccgit ggactigete egetgiegge alccagaaat tgegiggegg ageggeagae 780 gtgagccggc acggcaggcg gcctcctcct cctctcacgg cacggcagct acgggggatt 840 900 cetticecae egetecticg etillecette elegecegee giaalaaata gacaceceet ccacacccic iliccccaac cicgigiigi icggagcgca cacacacaca accagaicic 960

	ccccaaatcc	acccgtcggc	acctccgctt	caaggtacgc	cgctcgtcct	cccccccc	1020
_	ccctctctac	cttctctaga	tcggcgttcc	ggtccatggt	lagggcccgg	tagilclaci	1080
5	tetgtteatg	tttgtgttag	atccgtgttt	gigitagatc	cgtgctgcta	gcgttcgtac	1140
	acggatgcga	cctgtacgtc	agacacgitc	.tgattgctaa	cttgccagtg	tttctctttg	1200
10	gggaatcctg	ggatggctct	agccgttccg	cagacgggat	cgatttcatg	attititig	1260
	tttcgttgca	tagggtttgg	tttgcccttt	tcctttattt	caatatatgc	cgtgcacttg	1320
	titgtcgggt	catcitica	tgctttttt	tgtcttggtt	gtgatgatgt	ggtctggttg	1380
15	ggcggtcgtt	ctagaicgga	gtagaattct	gtttcaaact	acctggtgga	tttattaatt	1440
	tiggalcigi	atgtgtgtgc	catacatatt	catagitacg	aattgaagat	gatggatgga	1500
20	aatatcgatc	taggataggt	atacatgttg	atgcgggttt	tactgatgca	tatacagaga	1560
20	tgctttttgt	tcgcttggtt	gtgatgatgt	ggtgtggttg	ggcggtcgtt	cattegitet	1620
	agatcggagt	agaatactgt	ttcaaactac	ctggtgtatt	tattaatttt	ggaactgtat	1680
25	gtgtgtgtca	tacatcttca	tagttacgag	tttaagatgg	atggaaatat	cgatctagga	1740
	taggtataca	tgttgatgtg	ggttttactg	atgcatatac	atgatggcat	atgcagcatc	1800
	laticatatg	ctctaacctt	gagtacctat	ctattataat	aaacaagtat	gttttataat	1860
30	tattttgatc	ttgatatact	tggatgatgg	catatgcagc	agctatatgt	ggatttttt	1920
	agccctgcct	tcatacgcta	tttatttgct	tggtactgtt	tcttttgtcg	atgctcaccc	1980
35	tgttgtttgg	tgttacttct	gcaggtcgac	tctagaggat	ccgcgtccgg	gggcaatgag	2040
	ataigaaaaa	gcctgaactc	accgcgacgt	ctgtcgagaa	gttictgatc	gaaaagttcg	2100
	acagcgtctc	cgacctgatg	cagcicicgg	agggcgaaga	atctcgtgct	ttcagcttcg	2160
40	atgtaggagg	gcgtggatat	gtcctgcggg	taaatagctg	cgccgatggt	tictacaaag	2220
	alcgitatgt	ttatcggcac	tttgcatcgg	ccgcgctccc	gattccggaa	gtgcttgaca	2280
45	tiggggaati	cagcgagagc	cigacciaii	gcatcicccg	ccgtgcacag	ggtgtcacgt	2340
45	tgcaagacct	gcctgaaacc	gaactgcccg	ctgtictgca	gccggtcgcg	gaggccatgg	2400
	algcgatcgc	tgcggccgat	cttagccaga	cgagcgggtt	cggcccattc	ggaccgcaag	2460
50	gaatcggtca	atacactaca	tggcgtgait	tcatatgcgc	gattgctgat	ccccatgtgt	2520
	atcactggca	aacigigaig	gacgacaccg	tcagtgcgtc	cglcgcgcag	gctctcgatg	2580
	agcigaigci	ttgggccgag	gactgccccg	aagtccggca	ccicgigcac	gcggatttcg	2640
55	gctccaacaa	tglcctgacg	gacaalggcc	gcataacagc	ggtcattgac	tggagcgagg	2700

cgatgiicgg	ggattcccaa	tacgaggicg	ccaacatctt	ctictggagg	ccgtggttgg	2760
ctigiaigga	gcagcagacg	cgctacttcg	agcggaggca	tccggagctt	gcaggatcgc	2820
cgcggctccg	ggcgtatatg	ctccgcattg	gictigacca	actctatcag	agcttggttg	2880
acggcaattt	cgatgatgca	gcttgggcgc	agggtcgatg	cgacgcaatc	gtccgatccg	2940
gagccgggac	tgtcgggcgt	acacaaatcg	cccgcagaag	cgcggccgtc	tggaccgatg	3000
gctgtgtaga	agtactcgcc	gatagtggaa	accgacgccc	cagcactcgt	ccgagggcaa	3060
aggaatagag	tagatgccga	ccgaacaacg	gatccgtcga	cctgcagaga	tcgttcaaac	3120
attiggcaat	aaagtttctt	aagattgaat	cctgttgccg	gictigcgat	gattaicaia	3180
taatttctgt	tgaattacgt	taagcatgta	ataattaaca	tgtaatgcat	gacgilatii	3240
algagalggg	tttttatgat	tagagtcccg	caattataca	tttaatacgc	gatagaaaac	3300
aaaatatagc	gcgcaaacta	ggataaatta	tcgcgcgcgg	igicaiciat	gitaciagai	3360
cgatgataag	cigicaaaca	tgagaattca	gtacattaaa	aacglccgca	atgtgttatt	3420
aagttgtcta	agcgtcaatt	tgtttacacc	acaatatatc	ctgccaccag	ccagccaaca	3480
gcicccgac	cggcagcicg	gcacaaaatc	accactcgat.	acaggcagcc	catcagtccg	3540
ggacggcgtc	agcgggagag	ccgtigtaag	gcggcagact	ttgctcatgt	taccgatgct	3600
attcggaaga	acggcaacta	agctgccggg	tttgaaacac	ggatgatctc	gcggagggla	3660
gcatgttgat	tgtaacgatg	acagagcgtt	gctgcctgtg	atcaaatatc	atctccctcg	3720
cagagatccg	aattatcagc	cttcttattc	atttclcgct	taaccgtgac	aggctgtcga	3780
tcttgagaac	tatgccgaca	taataggaaa	tcgctggata	aagccgclga	ggaagctgag	3840
tggcgctatt	tctttagaag	tgaacgttga	cgatcgtcga	ccgtaccccg	atgaattaat	3900
tcggacgtac	gttctgaaca	cagciggata	cttacttggg	cgatigicat	acatgacatc	3960
aacaatgtac	ccgittgtgt	aaccgtctct	tggaggttcg	tatgacacta	giggiicccc	4020
tcagcttgcg	actagatgit	gaggaattcg	alctagtaac	atagatgaca	ccgcgcgcga	4080
taatttatcc	tagtttgcgc	gctatatttt	gtttictatc	gcgtattaaa	tgtataattg	4140
cgggactcta	atcataaaaa	cccatctcat	aaataacgic	atgcattaca	tgilaattat	4200
tacatgetta	acgtaattca	acagaaatta	tatgataatc	atcgcaagac	cggcaacagg	4260
attcaatctt	aagaaacttt	attgccaaat	giligaacga	tcaattcgag	ctcggtacgc	4320
aattccccac	cgaggctgta	gccgacgatg	gtgcgccagg	agagtigtig	aticatigii	4380
tgcctccctg	cigcggtttt	tcaccgaagt	tcatgccagt	ccagcgtttt	tgcagcagaa	4440

	aagccgccga	cticggiiig	cggtcgcgag	tgaagatccc	ttictigita	ccgccaacgc	4500
_	gcaatatgcc	ttgcgaggtc	gcaaaatcgg	cgaaattcca	laccigitca	ccgacgacgg	4560
5	cgclgacgcg	alcaaagacg	cggigalaca	tatccagcca	tgcacactga	tactcttcac	4620
	tccacaigic	ggtgtacatt	gagtgcagcc	cggclaacgt	atccacgccg	taticggiga	4680
10	tgataatcgg	ctgatgcagt	ttctcctgcc	aggccagaag	tictititce	agtacctict	4740
	ctgccgtttc	caaatcgccg	ctttggacat	accatccgta	ataacggttc	aggcacagca	4800
	catcaaagag	ategetgatg	gtatcggtgt	gagcgtcgca	gaacattaca	ttgacgcagg	4860
15	tgatcggacg	cgtcgggtcg	agtitacgcg	ttgcttccgc	cagiggcgaa	atattcccgt	4920
	gcacttgcgg	acgggtatcc	ggttcgttgg	caatactcca	catcaccacg	cttgggtggt	4980
20	ttttgtcacg	cgctatcagc	tctttaatcg	cctgtaagtg	cgcttgctga	giticcccgi	5040
	tgactgcctc	ttcgctgtac	agticiticg	gcttgttgcc	cgcttcgaaa	ccaatgccta	5100
	aagagaggtt	aaagccgaca	gcagcagiii	catcaatcac	cacgatgcca	tgitcaicig	5160
25	cccagtcgag	catctcttca	gcgtaagggt	aatgcgaggt	acggtaggag	ttggccccaa	5220
	iccagiccai	taatgcgtgg	tcgtgcacca	tcagcacgtt	atcgaatcct	ttgccacgta	5280
20	agtccgcatc	ttcatgacga	ccaaagccag	taaagtagaa	cggtttgtgg	ttaatcagga	5340
30	actgttcgcc	cttcactgcc	actgaccgga	tgccgacgcg	aagcgggtag	atatcacact	5400
	ctgtctggct	tttggctgtg	acgcacagt t	catagagata	accttcaccc	ggttgccaga	5460
35	ggtgcggatt	caccactigc	aaagtcccgc	tagtgccttg	tccagttgca	accaccigii	5520
	gateegeate	acgcagttca	acgctgacat	caccaitggc	caccacctgc	cagicaacag	5580
	acgcgtggtt	acagictigc	gcgacatgcg	tcaccacggt	galategice	acccaggtgt	5640
40	tcggcgtggt	gtagagcatt	acgctgcgat	ggaticcggc	atagttaaag	aaatcatgga	5700
	agtaagactg	ctttttcttg	ccgttttcgt	cggtaatcac	cattcccggc	gggatagtct	5760
45	gccagttcag	ttcgttgttc	acacaaacgg	tgatacgtac	actiticccg	gcaataacat	5820
	acggcgtgac	atcggcttca	aatggcgtat	agccgccctg	atgctccatc	acticcigat	5880
	tattgaccca	cactttgccg	taatgagtga	ccgcatcgaa	acgcagcacg	atacgctggc	5940
50	ctgcccaacc	tttcggtata	aagacticgc	gctgatacca	gacgitgccc	gcataattac	6000
	gaatatctgc	alcggcgaac	tgatcgttaa	aactgcctgg	cacagcaalt	gcccggcttt	6060
	cttgtaacgc	gctttcccac	caacgctgat	caattccaca	gitticgcga	tccagactga	6120
55	atgcccacag	gccgtcgagt	tttttgattt	cacgggttgg	ggtttctaca	ggacggacga	6180

	gtcgacggtt	cigiaaciai	catcatcatc	atagacacac	gaaataaagt	aatcagatia	6240
5	tcagttaaag	ctatgtaata	tttacaccat	aaccaatcaa	ttaaaaaata	gatcagttta	6300
•	aagaaagatc	aaagctcaaa	aaaataaaaa	gagaaaaggg	tcctaaccaa	gaaaatgaag	6360
	gagaaaaact	agaaatttac	cctgtaggga	tccatgttct	aggatctggt	tgtgtgtgtg	6420
10	tgcgctccga	acaacacgag	gttggggaaa	gagggtgtgg	agggggtgtc	tatttattac	6480
	ggcgggcgag	gaagggaaag	cgaaggagcg	gigggaaagg	aatccccgt	agctgccgtg	6540
	ccglgagagg	aggaggaggc	cgcctgccgt	gccggctcac	gtctgccgct	ccgccacgca	6600
15	attictggat	gccgacagcg	gagcaagtcc	aacggtggag	cggaactctc	gagaggggic	6660
	cagaggcagc	gacagagaig	ccgtgccgtc	tgcttcgctt	ggcccgacgc	gacgctgctg	6720
20	gttcgctggt	tggtgtccgt	tagactcgtc	gacggcgttt	aacaggctgg	cattatctac	6780
	tcgaaacaag	aaaaatgttt	ccttagtttt	tttaatttct	taaagggtat	iigiitaali	6840
	itiagicaci	ttattttatt	ciattttata	tctaaattat	taaataaaaa	aactaaaata	6900
25	gagititagi	tttcttaatt	tagaggctaa	aatagaataa	aatagatgta	ctaaaaaaat	6960
	tagictataa	aaaccattaa	ccctaaaccc	taaatggatg	tactaataaa	atggatgaag	7020
30	tattatatag	glgaagctal	ttgcaaaaaa	aaaggagaac	acatgcacac	taaaaagata	7080
	aaactgtaga	glcctgttgt	caaaatactc	aattgtcctt	tagaccaigt	ctaacigitc	7140
	atttatatga	-ttctctaaaa	cactgatatt	attgtagtac	tatagattat	attaticgia	7200
35	gagtaaagtt	taaatatatg	tataaagata	gataaactgc	acttcaaaca	agigigacaa	7260
	aaaaaatatg	tggtaatitt	ttataactta	gacatgcaat	gctcattatc	tclagagagg	7320
	ggcacgaccg	ggtcacgctg	cactgcaggc	algeaagete	ctaacatttt	attagagagc	7380
40	aggctagttg	cttagataca	tgatcttcag	gccgttatct	gtcagggcaa	gcgaaaatig	7440
	gccattlatg	acgaccaatg	cccgcagaa	gctcccatct	ttgccgccat	agacgccgcg	7500
45	ccccctttt	ggggtgtaga	acatectttt	gccagatgtg	gaaaagaagt	tcgitgiccc	7560
	attgttggca	atgacgtagt	agccggcgaa	agtgcgagac	ccatttgcgc	tatalataag	7620
	cctacgattt	ccgttgcgac	tattgtcgta	attggatgaa	ctattatcgt	agtigctcic	7680
50	agagttgtcg	taattigatg	gactattgtc	gtaattgctt	atggagttgt	cgtagttgct	7740
	tggagaaatg	tcgtagttgg	atggggagta	gtcataggga	agacgagcii	catccactaa	7800
56	aacaaltggc	aggicagcaa	gtgcctgccc	cgatgccatc	gcaagtacga	ggcllagaac	7860
55	caccttcaac	${\tt agatcgcgca}$	tagicticcc	cagcictcia	acgcligagi	taagccgcgc	7920

	cgcgaagcgg	cgtcggcttg	aacgaatigi	tagacattat	ttgccgacta	ccliggigat	7980
5	cicgcctitc	acgtagtgaa	caaattette	caactgatct	gcgcgcgagg	ccaagcgatc	8040
	ttcttgtcca	agataagcct	gcctagcttc	aaglatgacg	ggctgatact	gggccggcag	8100
	gcgctccatt	gcccagtcgg	cagcgacatc	cttcggcgcg	attttgccgg	ttactgcgct	8160
10	gtaccaaatg	cgggacaacg	taagcactac	atticgcica	tcgccagccc	agtcgggcgg	8220
	cgagttccat	agcgitaagg	titcatttag	cgcctcaaat	agatcctgtt	caggaaccgg	8280
	atcaaagagt	tcctccgccg	ciggacciac	caaggcaacg	clatgitete	tigcttitgt	8340
15	cagcaagata	gccagatcaa	tgicgatcgi	ggctggctcg	aagatacctg	caagaatgtc	8400
	attgcgctgc	cattetecaa	attgcagttc	gcgcttagct	ggataacgcc	acggaatgat	8460
20	gtcgtcgtgc	acaacaatgg	tgactictac	agcgcggaga	atctcgctct	ciccagggga	8520
	agccgaagtt	tccaaaaggi	cgttgatcaa	agctcgccgc	gtigtitcat	caagccttac	8580
	ggtcaccgta	accagcaaat	caatatcact	gtgtggcttc	aggccgccat	ccactgcgga	8640
25	gccgtacaaa	tgtacggcca	gcaacgtcgg	ttcgagatgg	cgctcgatga	cgccaactac	8700
	ctctgatagt	tgagicgata.	cttcggcgat	caccgcttcc	cicaigaigi	ttaactcctg	8760
30	aatlaagccg	cgccgcgaag	cggtgtcggc	tigaatgaat	tgttaggcgl	catcctgtgc	8820
	tcccgagaac	cagtaccagt	acatcgctgt	ticgticgag	acttgaggtc	tagittiata	8880
	cgigaacagg	tcaatgccgc	cgagagtaaa	gccacatttt	gcglacaaal	tgcaggcagg	8940
35	tacattgitc	gtitgigict	ctaatcgtat	gccaaggagc	tgtctgctta	gtgcccactt	9000
	tttcgcaaat	tcgatgagac	tgtgcgcgac	tcctttgcct	cggtgcgtgt	gcgacacaac	9060
	aatgigticg	atagaggeta	galcgitcca	tgttgagttg	agttcaatct	tcccgacaag	9120
40	cictiggicg	atgaatgcgc	catagcaagc	agagictica	tcagagicat	catccgagat	9180
	gtaatccttc	cggtaggggc	tcacacttct	ggtagatagt	tcaaagcctt	ggtcggatag	9240
45	gtgcacatcg	aacacttcac	gaacaatgaa	atggttctca	gcatccaatg	tttccgccac	9300
	cigcicaggg	atcaccgaaa	tcttcatatg	acgcctaacg	cciggcacag	cggatcgcaa	9360
	acciggcgcg	gcttttggca	caaaaggcgt	gacaggitig	cgaaiccgii	gcigccacit	9420
50	gttaaccctt	tigccagait	lggtaactal	aattlatgit	agaggcgaag	lcttgggtaa	9480
	aaactggcct	aaaattgctg	gggatttcag	gaaagtaaac	atcaccttcc	ggctcgatgt	9540
<i>55</i>	ctattglaga				_		9600
55	ggtgatgacg	gigaaaacci	ctgacacatg	cagctcccgg	agacggicac	agctigicig	9660

	laagcggatg	ccgggagcag	acaagcccgt	cagggcgcgt	cagcgggtgt	tggcgggtgl	9720
	cggggcgcag	ccaigaccca	gicacgiagc	gatagcggag	tgtatactgg	cttaactalg	9780
5	cggcatcaga	gcagattgla	ctgagagtgc	accatatgcg	gigigaaata	ccgcacagat	9840
	gcglaaggag	aaaataccgc	atcaggcgct	cttccgcttc	ctcgctcact	gactcgctgc	9900
10	gctcggtcgt	tcggctgcgg	cgagcggtat	cagcicacic	aaaggcggta	atacggttat	9960
	ccacagaatc	aggggataac	gcaggaaaga	acatgtgagc	aaaaggccag	caaaaggcca	10020
	ggaaccgtaa	aaaggccgcg	ttgctggcgt	ttttccatag	gctccgcccc	ccigacgagc	10080
15	atcacaaaaa	.tcgacgctca	agtcagaggt	ggcgaaaccc	gacaggacta	taaagatacc	10140
	aggcgtttcc	ccciggaagc	tccctcgtgc	gciciccigt	tccgaccctg	ccgcttaccg	10200
	gatacctgtc	cgcctttctc	ccttcgggaa	gcgtggcgct	ttctcatagc	tcacgctgta	10260
20	ggtatctcag	ttcggtgtag	gtcgttcgct	ccaagctggg	cigigigcac	gaaccccccg	10320
	ttcagcccga	ccgctgcgcc	ttatccggta	actatcgtct	tgagtccaac	ccggtaagac	10380
25	acgacttatc	gccactggca	gcagccactg	gtaacaggat	tagcagagcg	aggtatgtag	10440
	gcggtgctac	agagticitg	aagtggtggc	ctaactacgg	ctacactaga	aggacagtat	10500
	ttggtatctg	cgctctgctg	aagccagtta	ccttcggaaa	aagagttggt	agctcttgat	10560
30	ccggcaaaca	aaccaccgct	ggtagcggtg	gtttttttgt	ttgcaagcag	cagattacgc	10620
	gcagaaaaaa	aggatotoaa	gaagatcctt	tgatcttttc	tacggggtct	gacgctcagt	10680
35	ggaacgaaaa	cicacgiiaa	gggattitgg	tcatgagatt	atcaaaaagg	atcttcacct	10740
	agatectttt	aaattaaaaa	tgaagtttla	aatcaatcta	aagtatatat	gagtaaactt	10800
	•					tgictatiic	
40						gagggcttac	
	catctggccc	cagtgctgca	algataccgc	gagacccacg	ctcaccggct	ccagatttal	10980
						actitatecg	
45						ccagitaala	
						ttccattgtt	
50						cgctttcagc	
						ttatgacgaa	
						cgcgaggtcg	
55	ccgccccgta	accigicgga	tcaccggaaa	ggacccgtaa	agigataaig	attatcatct	11400

	acatatcaca	acgtgcgtgg	aggccatcaa	accacgicaa	ataatcaatt	atgacgcagg	11460
	tatcgtatta	attgatctgc	atcaacttaa	cgtaaaaaca	acttcagaca	atacaaatca	11520
5	gcgacactga	atacggggca	acctcatgtc	cccccccc	ccccctgc	aggcatcgtg	11580
	gtgtcacgct	cgtcgtttgg	tatggcttca	ttcagctccg	gttcccaacg	atcaaggcga	11640
o	gttacatgat	ccccatgit	gtgcaaaaaa	gcggttagct	ccttcggtcc	tccgatcgtt	11700
	gicagaagia	agttggccgc	agigitaica	ctcatggtta	tggcagcact	gcataattct	11760
	cttactgtca	tgccatccgt	aagatgcttt	tctgtgactg	gtgagtactc	aaccaagtca	11820
5	ttctgagaat	agtgtatgcg	gcgaccgagt	tgctcttgcc	cggcglcaac	acgggataat	11880
	accgcgccac	atagcagaac	tttaaaagtg	ctcatcattg	gaaaacgttc	ttcggggcga	11940
	aaactctcaa	ggatcttacc	gctgttgaga	tccagttcga	tgtaacccac.	tcgtgcaccc	12000
90	aactgatctt	cagcatcttt	tactttcacc	agcgtttctg	ggtgagcaaa	aacaggaagg	12060
	caaaatgccg	caaaaaaggg	aataagggcg	acacggaaat	gttgaatact	catactette	12120
5	ctttttcaat	attattgaag	cattlatcag	ggttattgtc	tcatgagcgg	atacatattt	12180
	gaatgtattt	agaaaaataa	acaaataggg	gttccgcgca	catttccccg	aaaagtgcca	12240
						gcgtatcacg	
10						ttttcgtgga	
						atcatcctgt	
5						acaagcagat	
						gctacgtccg	
						acccagacga	
0						tgggtggttg	
	•					giggiiggca	
						tecgitatic	
5						cactgatagt	
						gagtcacgit	
:o	atgacccccg	ccgatgacgc	gggacaagcc	gttttacgtt	tggaactgac	agaaccgcaa	
	cgttgaagga	gccactcagc	cc				12982

<210> 2

	<211> 90							
	<212> DNA							
5	<223>LB sequences are 1-25 and 55-70.							
	<400> 2							
10	ATTTACAATT GAATATATCC TGCCGccgct gccgctttgc acccgATTTA CAATTGAATA	60						
	TATCCTGCCG ccgctgccgc tttgcacccg	90						
15	<210> 3							
	<211> 90							
20	<212> DNA							
	<400> 3							
	cgggtgcaaa gcggcagcgg cggcaggata tattcaattg taaatcgggt gcaaagcggc	60						
25	agcggcggca ggatatattc aattgtaaat	90						
20	<210> 4							
30	<211> 25							
	<212> DNA							
35	<400> 4							
	catttctcgc ttaaccgtga caggc 25							
40	<210> 5							
	<211> 25							
45	<212> DNA							
	<400> 5							
	aaaccgcagc agggaggcaa acaat 25							
50								

Claims

- 1. A plant transformation vector based on the function of *Agrobacterium*, wherein the left border sequence has been modified such as to reduce the possibility of the integration of any non-T-DNA segment into plant chromosomes.
- 2. A plant transformation vector comprising a right border sequence and a left border sequence that can be recognized by the *vir* proteins of *Agrobacterium*, a T-DNA region located between these border sequences and into which a

gene to b introduced into the plant can be inserted, and a replication origin that nables replication of said vector in bacteria, wherein said left border sequence has been modified such as to reduce the possibility of integration of any non-T-DNA segment into plant chromosomes.

- 5 3. The plant transformation vector according to claim 1 or 2, wherein the modification of the left border sequence comprises more than one left border sequence.
 - 4. The plant transformation vector according to any one of claims 1 to 3, wherein the T-DNA sequence contains a marker gene that permits the selection of the transformant.
 - 5. The plant transformation vector according to any one of claims 1 to 4, wherein the replication origin permits replication of the vector in bacteria including bacteria for vector amplification and *Agrobacterium*.
- 6. A method of transforming plants comprising using an *Agrobacterium* host cell containing the vector according to any one of claims 1 to 5.
 - 7. A plant transformed by the method of claim 6.

10

20

25

30

35

40

45

50

55






Fig. 1



pSLB2



pSLB3




Fig. 2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/05386

	A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ Cl2N 15/82, A01H 5/00								
According to International Patent Classification (IPC) or to both national classification and IPC									
B. FIELD	DS SEARCHED								
	Minimum documentation searched (classification system followed by classification symbols) Int.Cl ² Cl2N 15/00-15/90								
L									
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched									
	data base consulted during the international search (nat SIS (DIALOG), WPI (DIALOG)	me of data base and, where practicable, sea	urch terms used)						
C. DOCU	MENTS CONSIDERED TO BE RELEVANT								
Category*	Citation of document, with indication, where a	ppropriate, of the relevant passages	Relevant to claim No.						
A	Plant Molecular Biology, 34, 1997 Allan Wenck et al., "Frequent collinear long transfer of DNA inclusive of the whole binary vector during Agrobacterium-mediated transformation" pp.913-922								
A	Proc. Natl. Acad. Sci. USA, 83, 1986 Horsch RB et al., "Rapid assay of foreign gene expression in leaf discs transformed by Agrobacterium tumefaciens: role of T-DNA borders in the transfer process" pp.4428-4432								
A	Plant Cell, 6, 1994 Martineau B. et al., "On defin pp.1032-1033	ing T-DNA"	1-7						
A	Proc. Natl. Acad. Sci. USA, 83 George C. Jen et al., "The right T-DNA is intrinsically more act region in promoting T-DNA trans pp.3895-3899	border region of pTiT37 ive than the left border	1-7						
Further	documents are listed in the continuation of Box C.	See patent family annex.							
"A" docume consider	categories of cited documents: Int defining the general state of the art which is not red to be of particular relevance document but published on or after the international filing	"T" later document published after the inte- priority date and not in conflict with th understand the principle or theory under "X" document of particular relevance; the c	e application but cited to crlying the invention						
"L" docume cited to	nt which may throw doubts on priority claim(s) or which is establish the publication date of another citation or other	considered novel or cannot be consider step when the document is taken alone "Y" document of particular relevance; the c	ed to involve an inventive laimed invention cannot be						
	rcason (as specified) nt referring to an oral disclosure, use, exhibition or other	considered to involve an inventive step combined with one or more other such combination being obvious to a person	documents, such						
"P" docume									
	ctual completion of the international search ecember, 1999 (17.12.99)	Date of mailing of the international scar 28 December, 1999 (2							
	ailing address of the ISA/ nese Patent Office	Authorized officer							
Facsimile No		Telephone No.							

Form PCT/ISA/210 (second sheet) (July 1992)



INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP99/05386

_			239703360
C (Continua	tion). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages		Relevant to claim No
A	Cell, 57, June 2, 1989 Guus Bakkeren et al, "Recovery of Agrobacterium tumefaciens T-DNA molecules from whole plants early after transfer", pp.847-857		1-7
A	Database BIOSIS on DIALOG, No.199900483906 Hanson Bill et al., "A simple method to enrich an Agrobacterium-transformed population for plants containing only T-DNA sequences", Plant Journal, 19(6), September, 1999, pp.727-734		1-7
A	Database BIOSIS on DIALOG, No.199598547959 Ramanathan V. et al., "Transfer of non-T-DNA p the Agrobacterium tumefaciens Ti plasmid pTiA left terminus of T-L-DNA", Plant Molecular Biolo 1995, pp.1149-1154	6 from the	1-7

Form PCT/ISA/210 (continuation of second sheet) (July 1992)